Mon. Not. R. Astron. Soc. **000**, 000–000 (2007) Printed 1 February 2008 (MN LaTeX style file v2.2)

arXiv:0709.0676v1 [astro-ph] 5 Sep 2007

The potential for Earth-mass planet formation around brown dwarfs

Matthew J. Payne[1] ⋆ and Giuseppe Lodato[2,1]
[1]*Institute of Astronomy, University of Cambridge, Madingley Road, Cambridge CB3 0HA*
[2]*Department of Physics and Astronomy, University of Leicester, Leicester, LE1 7RH*

Accepted —. Received —; in original form —

ABSTRACT

Recent observations point to the presence of structured dust grains in the discs surrounding young brown dwarfs, thus implying that the first stages of planet formation take place also in the sub-stellar regime. Here, we investigate the potential for planet formation around brown dwarfs and very low mass stars according to the sequential core accretion model of planet formation. We find that, for a brown dwarfs mass $0.05 M_{\odot}$, our models predict a maximum planetary mass of $\sim 5 M_{\oplus}$, orbiting with semi-major axis $\sim$ 1AU. However, we note that the predictions for the mass - semi-major axis distribution are strongly dependent upon the models chosen for the disc surface density profiles and the assumed distribution of disc masses. In particular, if brown dwarf disc masses are of the order of a few Jupiter masses, Earth-mass planets might be relatively frequent, while if typical disc masses are only a fraction of Jupiter mass, we predict that planet formation would be extremely rare in the substellar regime. As the observational constraints on disc profiles, mass dependencies and their distributions are poor in the brown dwarf regime, we advise caution in validating theoretical models only on stars similar to the Sun and emphasise the need for observational data on planetary systems around a wide range of stellar masses. We also find that, unlike the situation around solar-like stars, Type-II migration is totally absent from the planet formation process around brown dwarfs, suggesting that any future observations of planets around brown dwarfs would provide a direct measure of the role of other types of migration.

Key words: stars: low-mass, brown dwarfs – planetary systems: formation

1 INTRODUCTION

Observations of planets proceed apace: currently there are 232 extra solar planets known in 199 different planetary systems[1]. As is well known, many, if not all, of these systems are radically different to the Solar System (Beer et al. 2004), often posing challenges to theories of planet formation. As time has progressed, a number of planets have begun to be observed around low mass stars ($M_{\star} \sim 0.2 - 0.3 M_{\odot}$), thus begging the natural question of whether this trend will continue and whether similar planetary systems are to be expected at even lower masses, down in the brown dwarf regime.

Observations suggest that there does not seem to be any discontinuity between the properties of the discs around stars and those around brown dwarfs: Spitzer observations by Luhman et al. (2005) found that disc fractions for stars and brown dwarfs in the IC 348 and Chamaeleon I clusters were similar, thus suggesting a similar disc lifetime (see also Sterzik et al. 2004). The measurement by Scholz et al. (2006) of the Taurus star-forming region indicates that many of the discs around brown dwarfs must be relatively large in size, extending beyond 10AU, and have masses up to 12% of the central brown dwarf, with the average being 2.6%, compared with an average value of 1.9% for low mass main-sequence stars (note, however, that such disc mass estimates are very uncertain). Hence it appears that discs around brown dwarfs have similar properties and are as ubiquitous as discs around main sequence stars.

Observations indicate that grain growth, the precursor to planet formation in the sequential core accretion model, can occur efficiently in the small discs surrounding brown dwarfs (Apai et al. 2005). This then poses such questions as: what is the potential to form planets around very low mass stars; what would be their mass-semi-major axis distribution and would they be detectable? If formed or forming planets could be observed around brown dwarfs, these would provide highly valuable, low mass calibration points for the assessment of planet formation models over a wide range of stellar masses. In view of this, and in view of forthcoming missions such as the Terrestrial Planet Finder and Darwin, aimed at looking for Earth like planets, it is important to estimate the likelihood of the presence of planets around stars of different masses, and in particular around brown dwarfs.

Planet formation models currently fall into two categories: (i) the sequential core accretion model, which can account for the formation of both rocky planets and gas giants and (ii) the gravita-

---

⋆ E-mail: mpayne@ast.cam.ac.uk (MJP)

[1] http://exoplanet.eu, 06/05/07

tional instability model, which can only account for the formation of massive gas giants. The sequential core accretion model of planet formation (Pollack et al. 1996) assumes that planets form through a process in which first dust grains initially present in the protoplanetary disc condense to form kilometre-scale planetesimals; the collisional accumulation of planetesimals forms the basis for the runaway growth of a protoplanetary core of rock and/or ice (Safronov 1972; Wetherill 1980), and finally if a critical mass has been reached, the protoplanetary cores can begin to rapidly accrete a gaseous envelope (Mizuno 1980; Bodenheimer & Pollack 1986; Pollack et al. 1996). The gravitational instability model does not need to form the initial rocky core, and assumes that the protoplanetary disc may become gravitationally unstable and hence form gas-giants via gravitational collapse of high density regions in the disc (Boss 1998, 2000; Mayer et al. 2004). Whilst it is possible that some of the observed planets formed via a gravitational-instability scenario (and indeed also planetary mass companions to brown dwarfs. See, for example, the case of 2MASS1207B, Chauvin et al. 2005; Lodato et al. 2005), the results of Matsuo et al. (2007) suggest that around 90% of the observed extra-solar planets have a mass - semi-major axis relation consistent with the core-accretion model and that the metallicity dependence of the distribution of these planets can only be explained using the core-accretion model.

In a series of papers, Ida & Lin (2004a,b, 2005, henceforth IL1; IL2; IL3) attempted to construct a semi-analytic model of the combined core accretion and migration process which would allow quantitative comparison to be made between the observed distribution of extra-solar planets and that predicted by the theoretical core accretion model. They concentrated on investigating the observed mass - semi-major axis distribution of extrasolar planets, shown here in Fig. 1. This model starts with kilometre-size planetesimals, embedded in a two component gas and dust disc. The subsequent core accretion of planetesimals onto the protoplanet is modeled using the combined results of analytic models and N-body simulations. Once the cores reach a given critical mass, gas accretion is assumed to start, this process being modeled using numerical results for Kelvin-Helmholtz contraction of the gas envelope. This model also uses a number of prescriptions to enforce the limitation to core growth caused by various effects, such as isolation and migration.

Whilst this kind of modeling is admittedly oversimplified, the models of IL1-IL3 do succeed in reproducing several features of the observed distribution of extra-solar planets, such as the metallicity dependence for the probability of planet formation, the dependancy on stellar mass of the distribution of planetary masses as well as giving realistic mass ranges for the outcome of the planet formation process. However, the models experience problems in reproducing observations of intermediate mass planets with semi-major axes less than 3 AU. It is possible that these difficulties stem from the neglect of Type I migration in these models - see sections 2.4 and 3.2 for further consideration.

Clearly a self-consistent disc processing and core growth model for multiple interacting planets would be more appropriate (e.g. Laughlin et al. 2004; Papaloizou & Nelson 2005; Alibert et al. 2005; Papaloizou & Terquem 2006; Terquem & Papaloizou 2007). But given that the models of IL1-IL3 are currently the only extant attempt at providing a general framework for examining the outcome of the planetary formation and migration mechanism, they provide a useful framework for the exploration of the extensive parameter space and they allow us to start examining in a statistical manner the potential for planet formation in discs around brown dwarfs and the potential for practical observations in the near future.



Figure 1. Mass - Semi-major axis distribution of all currently known extra-solar planets.

To address the issue of planet formation around brown dwarfs under the core accretion model, we therefore extend the core-accretion model of IL1-IL3 down to the low mass regime. Once the basic structure of the model is in place, it can be used to investigate the effect on the planetary mass - semi-major axis distribution of varying key physical parameters. These include: the surface density profile of the disc, the size and mass of the disc, the rate of disc clearance, the effect of stellar mass and luminosity as well as migration.

2 PLANET FORMATION MODEL

The model we adopt here is primarily based on the model of sequential accretion presented in IL1 - IL3. In this section we initially briefly recount the principle components which pertain to our investigation before going on to discuss in more detail the refinements and additions which are required to extend the model down to the brown dwarf regime. Readers requiring more detail on the original model construction should refer to IL1 - IL3.

2.1 Disc Properties and Initial Conditions

The protoplanetary disc is modeled using a two component surface-density profile, $\Sigma = \Sigma_d + \Sigma_g$, to account for the dust and gas, with:

$$\Sigma_d = 10 f_0 \eta_{ice} \left(\frac{a}{1AU}\right)^k \left(\frac{M_\star}{M_\odot}\right)^n \text{g cm}^{-2} \qquad (1)$$

$$\Sigma_g = 2.4 f_0 \times 10^3 \left(\frac{a}{1AU}\right)^k \left(\frac{M_\star}{M_\odot}\right)^n e^{-t/\tau_{disc}} \text{g cm}^{-2}. \qquad (2)$$

where $M_\star$ is the mass of the central star, a is the radial distance from it, and η_{ice} is a step-function accounting for the increase in dust density at the snow line such that $\eta_{ice}(a < a_{ice}) = 1$ and $\eta_{ice}(a >$

$a_{ice}) = 4.2$. The radial location of the snow-line a_{ice} is set by the stellar luminosity, $L_\star$, such that (Hayashi 1981):

$$a_{ice} = 2.7\left(\frac{L_\star}{L_\odot}\right)^{1/2} \text{AU}. \tag{3}$$

The factors $(M_\star/M_\odot)^n$ allow an investigation of the dependence of the disc mass on stellar mass (here we examine the cases $n = 1$ and 2). The dissipation of the gaseous component is modeled as a simple exponential decay with characteristic timescale, τ_{disc}, chosen to be $\sim 10^6 - 10^7$ yrs, as suggested by observations of the infrared excess in young stellar objects (Haisch et al. 2001). The dust component is assumed to be time-independent. Setting the parameters $f_0 = 1$ and $k = -3/2$ would give the minimum mass nebula model (MMNM) (Hayashi 1981). We investigate the effect of varying the absolute surface density of the disc by varying f_0.

While IL1 - IL3 only consider the case where the surface density profile is a power law with index $k = -3/2$, here we also investigate the effects of having $k = -1$ (see section 3.4.1 for motivation and discussion).

It should also be noted that the simple exponential decay of the gaseous disc is the only evolution of the disc to be considered. The solid component is taken to be constant and neither component is modeled self-consistently with the growing mass of the planet. A self-consistent disc processing and core growth model would clearly be more appropriate (e.g. Alibert et al. 2005), but would also be more computationally demanding, so we adopt the simplified approach in this investigation.

To investigate the formation of planets around brown dwarfs, it is clear that we need to extend the semi-analytic model down to $M_\star < 0.084 M_\odot$. However, the model depends on $M_\star$ in two main ways: (i) Direct dependence on $M_\star$ via $\Sigma \propto M_\star^n$; and (ii) Indirect dependence on $M_\star$, e.g. the dependence of a_{ice} and hence Σ_d on $L_\star$. The detailed effects of varying $L_\star$ are discussed below in sections 2.3 and 3.4.3.

2.2 Core Growth and Gas Accretion

We follow the prescription for core growth and gas accretion as presented in IL1, with a core accretion timescale for a planet of mass M of (Kokubo & Ida 2002)

$$\begin{aligned}\tau_c &= 1.2\times10^5\left(\frac{\Sigma_d}{10\text{g cm}^{-2}}\right)^{-1}\left(\frac{a}{1AU}\right)^{3/5}\left(\frac{M}{M_\oplus}\right)^{1/3}\left(\frac{M_\star}{M_\odot}\right)^{-1/6} \\ &\quad\times\left(\frac{\Sigma_g}{2.4\times10^3\text{g cm}^{-2}}\right)^{-2/5}\left(\frac{m}{10^{18}g}\right)^{2/15} yr.\end{aligned} \tag{4}$$

where $m = 10^{18}$g is the assumed mass of the planetesimals accreted by the planetary core.

Ikoma et al. (2000) find that once a critical core mass of

$$M_{\text{crit}} = 10\left(\frac{\dot{M}}{10^{-6}M_\oplus\text{yr}^{-1}}\right)^{0.25} M_\oplus \tag{5}$$

has been reached, gas accretion can commence at a rate of

$$\tau_{K-H} = 10^{10}\left(\frac{M}{M_\oplus}\right)^{-3} yr. \tag{6}$$

Once a planet has grown to such as size that it opens a gap in the disc, the gas accretion rate is regulated by the rate at which the disc can process material and supply it across the gap to the planet. IL1 use a mass transfer rate given by the inferred rate for protostellar discs on T Tauri stars (Hartmann et al. 1998; Calvet et al. 2000)

$$\dot{M}_{disc} = 10^{-4}M_\oplus\left(\frac{t}{\tau_{disc}}\right)^{-3/2} \text{yr}^{-1}. \tag{7}$$

However, once the gap is opened, the accretion rate is reduced to values of the order of $= 0.1\times\dot{M}_{disc}$ (Artymowicz & Lubow 1996; Lubow & D'Angelo 2006). Here we have considered both the case described in Eq. (7), appropriately scaled down for brown dwarfs, and the case where accretion is further reduced due to gap opening, and found that this had little effect on the final distribution of planet masses and semi-major axes. Indeed, the details of gas accretion do not strongly affect our conclusions, since in most cases, as discussed below, we observe only very limited gas accretion in brown dwarf protoplanets.

We again acknowledge that these growth models are probably oversimplified. The form for τ_c given in Eq. (4) neglects the effects of core migration on the accretion rate of planetesimals onto the protoplanetary core (e.g. McNeil et al. 2005; Rice & Armitage 2003). The treatment of gas accretion giving rise to the expressions for M_{crit} and τ_{K-H} in Eqs. (5 and 6) is also highly approximate, neglecting the effects of opacity (Adams et al. 2005; Ikoma et al. 2000) and all the details evidenced by a numerical treatment of the problem (Bryden et al. 2000; Pollack et al. 1996).

2.3 Luminosity Evolution

The models of IL1 - IL3 treat the central star as a purely static object, with a fixed luminosity $L_\star$ equal to that on main-sequence, scaling such that $L_\star/L_\odot = (M_\star/M_\odot)^{1/4}$. For brown dwarfs, a constant luminosity model would be completely inappropriate, as no constant luminosity is ever reached, and we need to take this into account.

We allow stellar luminosity evolution to take place, accounting for this using the pre-main-sequence evolution simulations of Tout et al. (1999). Some sample evolutionary tracks are shown in Fig. 2. Compared to a static luminosity model, the net effect of the inclusion of luminosity evolution is to *raise* the luminosity at earlier times.

It should be noted that the core accretion model considered here, with an initial swarm of planetesimals of mass 10^{18}g, may have required $\sim 10^4$yrs for the planetesimals to form, although theoretical models vary considerably (Weidenschilling & Cuzzi 1993; Supulver et al. 1997; Rice et al. 2004, 2006). Hence we set the zero of time in our model to correspond to 10^4yrs in Fig.2.

2.4 Planetary Migration

Protoplanetary cores embedded within a gaseous disc interact with the gas and are thus subject to gravitational torques which cause them to migrate away from their initial position. For small core masses the interaction is linear and the corresponding migration regime is called Type I, while for larger core masses (typically above 1 Jupiter mass, for planets forming around a solar type star), a gap is opened in the disc and the resulting migration regime is called Type II (Ward 1997).

Type I migration is important for a number of reasons: (i) It is well known (Papaloizou & Terquem 2006) that the inclusion of Type-I migration in models of planet formation around solar-mass stars completely prevents the formation of massive gas giants (unless the disc mass is sufficiently small Thommes & Murray 2006), as all the planetary cores rapidly migrate into the central star, and



Figure 2. Luminosity evolution profiles from Tout et al. (1999) for (top-to-bottom) $M_\star/M_\odot$ = 1.5, 1.0, 0.6, 0.4, 0.2, 0.084 and 0.05 respectively.

thus Type I migration is generally neglected. However, simulations by Nelson & Papaloizou (2004) suggest that, in a turbulent disc, Type-I migration may actually follow a random walk, potentially providing a mechanism to ameliorate this problem. (ii) It may help to explain the presence of intermediate mass planets at small semi-major axes, by providing an additional migratory mechanism. In this paper, we have run models both including and neglecting the effect of standard (Ward 1997) Type-I migration.

Type-II migration occurs when the planet becomes massive enough to open a gap in the disc, the subsequent orbital evolution of the planet becoming firmly coupled to the viscous evolution of the disc (Lin & Papaloizou 1985). A gap is opened up when the planetary mass exceeds

$$\begin{aligned}\frac{40\nu}{a^2\Omega_K}M_\star &= 40\alpha\left(\frac{c_s}{V_K}\right)^2 M_\star \\ &= 3\left(\frac{\alpha}{10^{-4}}\right)\left(\frac{a}{1AU}\right)^{1/2}\left(\frac{L_\star}{L_\odot}\right)^{1/4} M_\oplus, \qquad (8)\end{aligned}$$

where we have used the α-parameterization for the effective viscosity, $\nu = \alpha H^2\Omega_K$, of Shakura & Sunyaev (1973) and we use $c_s^2 \propto T$.

However, hydrodynamical simulations (Lin & Papaloizou 1985; Ida & Lin 2004a) suggest that the onset of Type-II migration occurs somewhat later, when the planet has grown to a larger mass. Type-II migration is therefore taken to begin when the planet exceeds 10 times the gap opening mass.

In general, in any viscously evolving gaseous disc, the inner parts move in and accrete, while the outer parts move out and take up the angular momentum lost by the accretion material (Lynden-Bell & Pringle 1974). The direction of Type II migration is therefore determined by the position of the planetary core with respect to the radius which marks the transition from the inner to the outer disc, which we obtain from the appropriate self-similar solution of Lynden-Bell & Pringle (1974).

2.5 Methodology

Given the above semi-analytic prescription for protoplanetary growth, if we specify $M_\star$, k, n, f_0, τ_{disc} and the initial semi-major axis a_i, we can then allow the planet to grow from an initial mass of 10^{22} g. If we then perform a Monte-Carlo simulation, allowing the parameters f_0, τ_{disc} and a_i to vary, we can build up a picture of the potential for planet formation for a wide range of protoplanetary discs around a wide range of stars.

Because the initial size distribution is unknown for brown dwarf disks, the dissipation timescale is also unknown. Attempts at estimating this timescale have been made by Alexander & Armitage (2006) based on the $\dot{M} - M$ relation (Natta et al. 2004), and they argue that $t \sim M_\star^{-1}$. However, it is not clear whether such observational scalings really reflect some intrinsic properties of the initial protostellar disc population, nor how much are they affected by selection effects (Clarke & Pringle 2006), so we have kept the same timescale for solar mass stars and brown dwarfs (but we do allow for a relatively large spread in timescale, as suggested by Clarke (2007)). We therefore allow τ_{disc} to be evenly distributed in logarithmic scale over $10^6 - 10^7$yrs.

We take f_0 to be log-normally distributed (centred on $\log f_0 = 0$ and with variance equal to 1) and then demand that the disc mass within 100AU be less than $0.35M_\star$, to ensure that the disc is gravitationally stable out to 100AU. We thus end up with a distribution of disc masses such that the maximum disc/stellar mass fraction is $\leqslant 0.35$, with an average significantly below this value. Note that Rodríguez et al. (2005) have observed protostellar disc masses as high as $0.35M_\star$, indicating that the above upper limit is reasonable. Finally, we distribute the initial orbital semi-major axes evenly in log scale over 0.1 to 100 AU and terminate all migration at 0.04AU[2]

3 RESULTS

Before discussing our results on planet formation around brown dwarfs, we initially test our model by reproducing some of the main results of IL1 - IL3 for planets around Solar-type stars. Our standard model assumes a disc profile with $\Sigma \propto a^{-3/2}$ and $f_0 \propto (M_\star/M_\odot)^2$ and considers only Type-II migration. We then extend our standard model down to the Brown Dwarf regime, adding in also the effect of luminosity evolution (section 3.3). Finally, we investigate the impact of varying our main parameters on the potential for planet formation (section 3.4).

3.1 Planet Formation around Solar-Mass Stars

Fixing $M_\star = M_\odot$, we reproduce in Fig. 3 the Mass - Semi-Major axis distribution as found in IL1, fig. 12c. The model uses our standard values such that $k = -3/2$ in $\Sigma \propto (a/1\mathrm{AU})^k$ and the distribution of values of f_0 is such that the maximum disc mass is $0.35M_\star$, with an average of $0.05M_\star$. We use the following nomenclature, approximately equivalent to that of IL1, to label the planets:

- Gaseous: $M_g > 10M_c$;
- Intermediate: $M_c < M_g < 10M_c$;

---

[2] While in general, for solar-type stars, this radius corresponds to the stellar magnetospheric radius, there is no strong reason for assuming that the termination radius in brown-dwarf discs would be the same, but as we show later, we find little or no migration around brown dwarfs, so its exact value is effectively unimportant.



Figure 3. Planet masses $> 1M_\oplus$ for $M_\star = M_\odot$ with static luminosity. The different symbols respectively label: Open Circles - "Gaseous" ($M_g > 10M_c$); Filled Triangles - "Intermediate Composition" ($10M_c > M_g > M_c$); Crosses - "Rocky" ($M_g < M_c$).



Figure 4. Planet masses $> 1M_\oplus$ for $M_\star = 0.3M_\odot$ with only Type II migration present. Labels are as in Fig. 3, with the addition of observed planets plotted as filled squares.

- Rocky: $M_g < M_c$;

where M_g is the final gas mass and M_c is the core mass, such that the total planet mass is $M = M_g + M_c$.

The model distribution presented in Fig. 3 faithfully reproduces that of IL1, fig. 12c: They both display the characteristic "planet desert" for $a < 4$AU and $10M_\oplus < M < 100M_\oplus$ and both have a maximum gas-giant mass of $\sim 3000M_\oplus$. In addition, the distribution of planet types within the diagram is very similar (although the precise definition of planet-type-labels may differ slightly).

Note the high density of massive planets at $a = 0.04$AU caused by the inward Type-II migration being arbitrarily halted at this distance.

Note also that below the $1M_\oplus$ line, there is a vast tail of small-mass planets produced by the simulations; planets which grew only very slowly and/or were growing in a disc whose mass was very small.

3.2 Planet Formation around Intermediate-Mass Stars

The results of Fig 3 are in good general agreement with observed extra-solar planets, the exception being the problematic observations of intermediate mass planets at small semi-major axes. IL3 note that their results indicate that close-in, intermediate mass planets cannot form around F,G and K dwarfs, but may form around M dwarfs and note that the formation of dynamically isolated low mass planets around such stars would likely be attributable to Type I migration or the effect of sweeping secular or mean-motion resonances.

In Fig 4 we focus on the observed planetary systems around Gl-581 and Gliese 876, both of which have $M_\star \sim 0.3M_\odot$. We thus plot the observed planets on the same scale as our numerical results, using models that do not include Type I migration. We consistently find it very difficult to produce close in, intermediate mass planets (such as Gl-581-c Udry et al. 2007), if we do not include some form of Type I migration in the process.

However, as mentioned above, the simple introduction of the fast Type I migration (Ward 1997) would generally preclude the formation of the high mass ($\sim 100M_\oplus$) planets observed in these systems, unless the migration rate is significantly slower than the standard analytic form (Nelson & Papaloizou 2004; Thommes & Murray 2006). It thus appear that, while some additional form of planetary core migration is needed in order to reproduce the above-mentioned observations, its rate should be substantially reduced with respect to the simple analytical estimates. In the rest of the paper we will therefore consider the two extreme cases in which we neglect completely Type I migration and in which we include standard Type I migration.

3.3 Planet Formation around Brown Dwarfs

If we now allow the central stellar mass $M_\star$ to vary down below $0.084M_\odot$, we can to investigate the potential for planet formation in the brown dwarf regime. As noted in section 2.3, this extension requires the implementation of a luminosity evolution model for brown dwarfs.

In Fig. 5 we plot the mass - semi-major axis distribution resulting from our numerical simulation of the planet formation process around brown dwarfs of mass $0.05M_\odot$ - note the change in scale compared to Fig. 3. We keep the standard disc profile of section 3.1 and assume that the disc surface density scales as $M_\star^2$ (i.e., we assume $n = 2$ in Equation (2)). This effectively makes the $0.35M_\star$ cut-off unnecessary, as the maximum disc mass is now $0.21M_\star$ (which corresponds to $10.5M_{\rm Jupiter}$), while the average disc mass is $0.003M_\star$ ($0.15M_{\rm Jupiter}$).

Clearly the outcome of the planet formation process around brown dwarfs is radically different to that for G-dwarfs. Of particular importance is the fact that all of the planets produced are of



Figure 5. Planet Masses for Brown Dwarf with $M_\star = 0.05M_\odot$. Labels are as in Fig. 3.



Figure 6. Planet Masses $> 1M_\oplus$ for $M_\star = M_\odot$ with $k = -1$. Labels are as in Fig. 3.

low mass - we do not see any significant number of planets formed above $1M_\oplus$, the most massive ones corresponding to the high tail of the disc mass distribution - and are of an icy constitution. In none of the simulation runs did any planet grow massive enough for any significant gas accretion take place. This means that the distribution of planetary masses is effectively limited by the surface density of the solid disc. The radial extent of the planet population is at least a factor of ten smaller than around solar-mass stars and we find that the peak of the distribution shifts to significantly smaller radii as the stellar mass decreases, reflecting the fact that disc densities at any given radii are lower around brown dwarfs.

Whilst the lack of gas accretion limits the predicted mass of the planets, it also has implications for planetary migration. If a planet does not accrete gas and remains relatively low mass, it is highly unlikely to open up a gap in the gaseous disc, and hence will not under-go Type-II migration. This means that the only migration mechanism relevant for the brown dwarf regime will be Type-I (which has not been included here). Thus, any observations of planets around brown dwarfs could shed light directly on the effects of Type-I migration and hence illuminate its likely character in planet formation around all higher mass stars.

Note also the paucity of planets plotted in Fig. 5, indicating that Earth-like planet formation is highly unlikely in this case. Indeed, we found a planet with mass $M > 0.3M_\oplus$ in only 0.035% of the simulations that we have performed.

3.4 Impact of Parameter Variation on Planet Formation

3.4.1 Surface Density Dependence on Semi-Major Axis

It was assumed in section 3.3 that the disc profile varied as $\Sigma \propto a^{-3/2}$. In this section we explore the effect of having a shallower density profile, with $\Sigma \propto a^{-1}$, while keeping the scaling with stellar mass as $(M_\star/M_\odot)^2$, and allowing only Type II migration. We find that at both solar mass and lower masses of $M_\star = 0.05M_\odot$, the effect of moving to $k = -1$ is significant.

At $M_\star = M_\odot$, moving to a $k = -1$ model leaves the average and maximum *disc* masses unchanged from the standard model in section 3.1. However, a comparison between Fig. 6 (which shows the planet mass - semi-major axis distribution in this case) and Fig. 3 shows that for semi-major axes $<$ 10AU, the change in model gives very similar results for the distribution of Jupiter-mass planets, but suppresses the production of terrestrial planets. Fig. 7 shows the average and maximum planetary masses at different radii for the various cases. This shows that both the maximum and average planet mass inside 10AU (which are dominated by the gas giants) are similar for these models, but beyond 10AU, the effect of the $k = -1$ model is to increase both the average and maximum plant masses. This result can be simply understood based on the fact that, for a given disc mass, with a shallower surface density profile the disc density is decreased at small radii and increased at large radii. Note that Raymond et al. (2005) looked at terrestrial planet formation at small radii and found it was suppressed in discs with shallower profiles, in agreement with our result.

Let us now turn to analyse the case of brown dwarf. Fig. 8 shows the planet mass - semi-major axis distribution for $k = -1$ (cf. Fig. 5). The effect of moving to a $k = -1$ model, similarly to the case of higher stellar mass, is to suppress the formation of rocky planets. This is further evidenced in Fig. 9, which shows a comparison of the maximum and average planet mass at different radii for the two cases $k = -1$ (solid line) and $k = -3/2$ (dashed line). Unlike the case of Solar-type stars, these histograms are now dominated by rocky planets, and indeed moving to a shallower surface density profiles results in a reduction of both the maximum and average mass planet formed, preventing the formation of any planets with masses greater than $1M_\oplus$.

3.4.2 Surface Density Dependence on Stellar Mass

Disc masses estimates for brown dwarfs are very uncertain and only known for a handful of objects (Scholz et al. 2006). In our standard model, (section 3.3) we have assumed that protostellar disc masses



Figure 7. Changing the slope of the surface density profile at $M_\star = 1M_\odot$. The thin lines indicate *average* masses, the thick *maximum* masses, whilst solid lines indicate $k = -1$, dotted lines $k = -3/2$.



Figure 9. Changing the slope of the surface density profile at $M_\star = 0.05M_\odot$. The thin lines indicate average masses, the thick maximum masses, whilst solid lines indicate $k = -1$, dotted lines $k = -3/2$.



Figure 8. Planet Masses versus semi-major axis for $M_\star = 0.05M_\odot$ with $k = -1$. Labels are as in Fig. 3.

scale with stellar mass as $M_\star^2$, leading to typical disc masses of a tenth of a Jupiter mass in the brown dwarf regime. Here, we examine the effect of allowing for relatively more massive discs around brown dwarfs, and we thus consider the case where the disc surface density scales linearly with stellar mass.

We plot in Fig. 10 the resulting planet mass versus semi-major axis distribution in this case, for the two cases where $k = -3/2$ (steeper density profile; left panel) and $k = -1$ (shallower density profile; right panel) and collate in Fig. 11 the histograms of average and maximum planet masses formed around brown dwarfs for the various models.

In this case, typical disc masses are of the order of a few Jupiter masses and the enhanced disc mass is reflected in a larger fraction of objects being able to form Earth-like planets. Indeed, planets with mass larger than $0.3M_\oplus$ are formed now in ≈10% of the cases for $k = -3/2$ and in ≈ 2% of the cases for $k = -1$.

In this case, the cores are growing faster, but still none of them grow quickly enough to start accreting gas before it dissipates. Thus, despite the increased potential for planet formation, no object is able to accrete a significant gaseous envelope, and the maximum planet mass is only slightly enhanced, the maximum now being $5M_\oplus$, in the most favourable case (that is, $k = -3/2$).

The maximum planetary mass obtained is also slightly dependent on the upper limit on the disc mass in order for the disc to be gravitationally stable. For example, if we only allowed discs less massive than 10% of the central brown dwarf mass, the maximum resultant planetary mass would be decreased to 1 $M_\oplus$ and it would be raised to $9M_\oplus$, with some very limited gas accretion occurring, if the discs are allowed to be half as massive as the brown dwarf.

Clearly, an important parameter in the determination of the final planet population is the disc surface density. For a given total disc mass, a more compact disc, with an outer radius significantly smaller than the 100 AU assumed above, would have a larger surface density and therefore an enhanced chance of forming planets. We have thus re-run the most favourable case for planet formation (that is, the one with $n = 1$ and $k = -3/2$), assuming the same distribution of total disc mass, but with significantly smaller disc sizes (this, in turn, implies increasing the factor f_0 in Eqs. (1) and (2)). The results are shown in Fig. 12, for an outer disc radius $R_{\rm out} = 50$ AU (left panel) and 20 AU (right panel). While the reduction of $R_{\rm out}$ to 50 AU only results in a slightly increased maximum planet mass, in the case of $R_{\rm out} = 20$ AU we do see the appearance of a number of gas giants in a limited number of cases. However, we regard these cases as extremely unlikely, since they correspond to



Figure 10. Planet Masses versus semi-major axis for Brown Dwarf with $M_\star = 0.05 M_\odot$, $n = 1$ and $k = -3/2$ (left panel) and $k = -1$ (right panel). Labels are as in Fig. 3.



Figure 11. Results for planet formation around brown dwarfs. The left panel shows the average planet masses while the right panel shows the maximum planet mass in the various models considered. Solid Line - $n = 1$ and $k = -3/2$; Dotted Line - $n = 1$ and $k = -1$; Dashed Line - $n = 2$ and $k = -3/2$; Dot/Dash Line - $n = 2$ and $k = -1$.

the upper tail of the disc mass distribution, with disc masses of the order of several Jupiter masses, confined within a region as small as 20 AU.

3.4.3 Introduction of luminosity evolution

As mentioned above, the models that we use for planet formation around brown dwarfs also include the effect of the evolution of the (sub)-stellar luminosity with time. The stellar luminosity has two main effects on our models: (a) it determines the temperature of the gas and (b) it sets the location of the snow-line, inside which



Figure 12. Planet mass versus semi-major axis for $M_\star = 0.05M_\odot$, $n = 1$ and $k = -3/2$, with outer disc radius equal to 50 AU (left panel) and 20 AU (right panel). While in the case of $R_{\rm out} = 50$ AU, the final distribution of planet mass is very similar to the case of $R_{\rm out} = 100$ AU, some gas giants appear in the case $R_{\rm out} = 20$ AU. These latter cases, however, correspond to the very unlikely case of a massive disc ($M_{\rm disc} \gtrsim$ a few Jupiter masses) with a very small outer radius.

the solid density is significantly reduced (we neglect here any other possible effect related to changes in the efficiency of angular momentum transport within the discs). Since there is little or no gas accretion on proto-planets around brown dwarfs in our model, and no planet becomes massive enough to open up a gap and start migrating, the first of the above effects actually has essentially no impact on the final outcome. The second effect is indeed present, but it turns out not to affect deeply the final results, and simulations performed without taking into account the luminosity evolution of the brown dwarfs only show relatively small differences.

3.4.4 Introduction of planetary migration

As IL3 have already noted, since low-mass cores do not undergo Type II migration, any detection of close-in super-Earths around low mass stars, as in the case of Gl-581-c (a $5.5M_\oplus$ planet orbiting a $0.3M_\odot$ star, Udry et al. 2007), would need the inclusion of other types if migration. In our models of planetary formation around brown dwarfs in which Type I migration is neglected, we always predict a peak in the distribution at 1AU, irrespective of the other parameter variations in the model.

As discussed in section 3.2, we now consider the addition of a form of Type I migration to our model. In particular, we consider the model for Type-I migration as derived by Tanaka et al. (2002) for a 3-dimensional isothermal gaseous disc, where the migration rate is of the form

$$\tau_I \approx 1\times10^5 \left(\frac{M}{M_\oplus}\right)^{-1} \left(\frac{M_\star}{M_\odot}\right)^{3/2} \left(\frac{\Sigma_g}{2\times10^3 {\rm g\ cm}^{-2}}\right)^{-1} {\rm yrs}, \qquad (9)$$

We find that in general the effects of Type I migration around brown dwarfs are less significant than around solar type stars. This is mostly due to the fact that planetary cores around brown dwarfs grow at a much slower rate, and therefore the migration timescale in this case is relatively longer. For the $n = 1$ $k = -3/2$ model (see Fig. 13), the maximum planet mass at $\sim$ 1AU is very slightly reduced, whilst the average value at this radius (and the efficiency of the planet formation process) is reduced as a large number of the cores are swept into the inner boundary.

Since Type-II migration effectively never occurs around brown dwarfs, any observational evidence of a secondary peak/excess in the radial distribution of planetary semi-major axes inside the main distribution peak at $\sim$ 1AU would provide a clean, direct insight into the nature and extent of Type-I migration.

4 DISCUSSION AND CONCLUSIONS

We have extended standard core-accretion models for planet formation down to the brown dwarf regime and investigated the potential for planet formation in protoplanetary discs around objects with masses $\sim 0.05M_\odot$. We looked at the impact on the predicted planetary mass - semi-major distribution of varying fundamental parameters, like the total disc mass and its radial distribution.

In line with previous results (IL1), at higher stellar masses ($M_* \sim 1M_\odot$, Fig. 3), we found that the full range of extant planet types is produced, including rocky and icy cores up to $10^2M_\oplus$, rock and ice giants spanning a range $10^1 - 10^3M_\oplus$ and gas giants extending from $10^2M_\oplus$ up to $10^4M_\oplus$. In addition, we have found that a shallower surface density profile results in a reduced efficiency of the formation of rocky planets at radii $\lesssim$ 10 AU.

Our main result is the determination of the likelihood of planet formation around brown dwarfs, with mass $\sim 0.05M_\odot$. Our main findings can be summarized as follows. (*i*) Giant planet formation is completely inhibited in this case. In none of our standard simulations did any planet accrete a significant gaseous envelope. This occurs despite the fact that in a few cases relatively massive



Figure 13. Effect of analytic Type I migration on brown dwarfs with $M_\star = 0.05M_\odot$ and $n = 1$, $k = -3/2$. Labels are as in Fig. 3.

cores do form (in principle massive enough to start runaway gas accretion), because such massive cores, when they form, only do so at a late time, when most of the gaseous component of the disc has dissipated. This means that Jupiter mass companions to brown dwarfs, as sometimes observed, have to be formed through a different – binary like – mechanism, that is either core or disc fragmentation. While this had already been discussed for very wide extremely low mass binaries, like 2MASS1207 (Lodato et al. 2005), we can now extend this conclusion to closer binaries. (*ii*) The formation of Earth-like planets is possible even around brown dwarfs, and the maximum planet mass that we have found is $\approx 5M_\oplus$. However the likelihood of this process is critically dependent on what the typical disc masses are in this regime. In particular, if typical brown dwarf disc masses are of the order of a few Jupiter masses (which is consistent with a linear scaling of disc mass with stellar mass), then up to 10% of brown dwarfs might possess planets with masses $> 0.3M_\oplus$, while on the other hand if typical brown dwarf disc masses are only a fraction of a Jupiter mass (consistent with a quadratic scaling of disc mass with stellar mass), then even the formation of Earth-like planets would be essentially prohibited. We note that current estimates of brown dwarfs disc masses are only available for a small number of objects and are very close to our boundary between planet-forming and non-planet-forming systems, with upper limits at the level of a few Jupiter mass (Scholz et al. 2006). In order to answer the question of the occurrence of planets around brown dwarfs it is therefore essential to improve the statistics and the accuracy of disc mass measurements in the very low mass regime.

Finally, we note that the lack of any gas accretion in the planet formation process around brown dwarfs and the concomitant lack of any Type-II migration means that the radial distribution of the planets will be determined purely by the initial mass distribution in the disc and the subsequent effects of Type-I migration. Given that the underlying disc models tend to produce the most massive planets at $a \sim 1$AU, irrespective of the specific model, then any excess/secondary peak (or lack thereof) in the number of detected planets inside this radius would give clear evidence of the nature of Type-I migration.

ACKNOWLEDGMENTS

We would like to thank Daniel Apai, Cathie Clarke, Ilaria Pascucci, Doug Lin, Jim Pringle and Chris Tout for helpful discussions. MJP acknowledges the UK PPARC/STFC for a research studentship.

REFERENCES

Adams F. C., Bodenheimer P., Laughlin G., 2005, Astronomische Nachrichten, 326, 913
Alexander R. D., Armitage P. J., 2006, ApJ, 639, L83
Alibert Y., Mordasini C., Benz W., Winisdoerffer C., 2005, A&A, 434, 343
Apai D., Pascucci I., Bouwman J., Natta A., Henning T., Dullemond C. P., 2005, Science, 310, 834
Artymowicz P., Lubow S. H., 1996, ApJ, 467, L77+
Beer M. E., King A. R., Livio M., Pringle J. E., 2004, MNRAS, 354, 763
Bodenheimer P., Pollack J. B., 1986, Icarus, 67, 391
Boss A. P., 1998, Nat, 393, 141
Boss A. P., 2000, ApJ, 536, L101
Bryden G., Lin D. N. C., Ida S., 2000, ApJ, 544, 481
Calvet N., Hartmann L., Strom S. E., 2000, Protostars and Planets IV, pp 377
Chauvin G., Lagrange A.-M., Dumas C., Zuckerman B., Mouillet D., Song I., Beuzit J.-L., Lowrance P., 2005, A&A, 438, L25
Clarke C. J., 2007, MNRAS, 376, 1350
Clarke C. J., Pringle J. E., 2006, MNRAS, 370, L10
Haisch Jr. K. E., Lada E. A., Lada C. J., 2001, ApJ, 553, L153
Hartmann L., Calvet N., Gullbring E., D'Alessio P., 1998, ApJ, 495, 385
Hayashi C., 1981, Progress of Theoretical Physics Supplement, 70, 35
Ida S., Lin D. N. C., 2004a, ApJ, 604, 388 (IL1)
Ida S., Lin D. N. C., 2004b, ApJ, 616, 567 (IL2)
Ida S., Lin D. N. C., 2005, ApJ, 626, 1045 (IL3)
Ikoma M., Nakazawa K., Emori H., 2000, ApJ, 537, 1013
Kokubo E., Ida S., 2002, ApJ, 581, 666
Laughlin G., Bodenheimer P., Adams F. C., 2004, ApJ, 612, L73
Lin D. N. C., Papaloizou J., 1985, in Black D. C., Matthews M. S., eds, Protostars and Planets II. p. 981
Lodato G., Delgado-Donate E., Clarke C. J., 2005, MNRAS, 364, L91
Lubow S. H., D'Angelo G., 2006, ApJ, 641, 526
Luhman K. L. et al., 2005, ApJ, 631, L69
Lynden-Bell D., Pringle J. E., 1974, MNRAS, 168, 603
Matsuo T., Shibai H., Ootsubo T., Tamura M., 2007, astro-ph/0703237
Mayer L., Quinn T., Wadsley J., Stadel J., 2004, ApJ, 609, 1045
McNeil D., Duncan M., Levison H. F., 2005, AJ, 130, 2884
Mizuno H., 1980, Progress of Theoretical Physics, 64, 544
Natta A., Testi L., Muzerolle J., Randich S., Comerón F.,Persi P., 2004, A&A, 424, 603
Nelson R. P., Papaloizou J. C. B., 2004, MNRAS, 350, 849
Papaloizou J. C. B., Nelson R. P., 2005, A&A, 433, 247
Papaloizou J. C. B., Terquem C., 2006, Reports of Progress in Physics, 69, 119
Pollack J. B., Hubickyj O., Bodenheimer P., Lissauer J. J., Podolak M., Greenzweig Y., 1996, Icarus, 124, 62
Raymond S. N., Quinn T., Lunine J. I., 2005, ApJ, 632, 670
Rice W. K. M., Armitage P. J., 2003, ApJ, 598, L55
Rice W. K. M., Lodato G., Pringle J. E., Armitage P. J., Bonnell I. A., 2004, MNRAS, 355, 543
Rice W. K. M., Lodato G., Pringle J. E., Armitage P. J., Bonnell I. A., 2006, MNRAS, 372, L9
Rodríguez L. F., Loinard L., D'Alessio P., Wilner D. J., Ho P. T. P., 2005, ApJ, 621, L133

Safronov V. S., 1972. Evolution of the protoplanetary cloud and formation of the earth and the planets. Jerusalem

Scholz A., Jayawardhana R., Wood K., 2006, ApJ, 645, 1498

Shakura N. I., Sunyaev R. A., 1973, A&A, 24, 337

Sterzik M. F., Pascucci I., Apai D., van der Bliek N., Dullemond C. P., 2004, A&A, 427, 245

Supulver K. D., Bridges F. G., Tiscareno S., Lievore J., Lin D. N. C., 1997, Icarus, 129, 539

Tanaka H., Takeuchi T., Ward W. R., 2002, ApJ, 565, 1257

Terquem C., Papaloizou J. C. B., 2007, ApJ, 654, 1110

Thommes E. W., Murray N., 2006, ApJ, 644, 1214

Tout C. A., Livio M., Bonnell I. A., 1999, MNRAS, 310, 360

Udry S. et al., 2007, astro-ph/0704.3841

Ward W. R., 1997, Icarus, 126, 261

Weidenschilling S. J., Cuzzi J. N., 1993, in Levy E. H., Lunine J. I., eds, Protostars and Planets III. p. 1031

Wetherill G. W., 1980, ARA&A, 18, 77